The Bryologist 97(4), 1994, pp. 476–486
Copyright © 1994 by the American Bryological and Lichenological Society, Inc.

Index to Volume 97 of THE BRYOLOGIST[1]

PIERS MAJESTYK
Biology Department, University of Southwestern Louisiana, Lafayette, LA 70504-2451

Abrothallus parmeliarum, 156+
Acacia mearnsii, 298+
Acarospora chlorophana, 142+; fuscata, 142+; glaucocarpa, 142+; gwynni, 440+; schleicheri, 142+; smaragdula, 142+; superfusa, 142+; veronensis, 142+
Acrocryphaea, 89+
Acromastigum colensoana, 319+; cunninghamii, 319+*; mooreanum, 314
Acroscyphus sphaerophoroides, 56+
Actinogyra muehlenbergii, 142+
Actinothuidium hookeri, 128+
Adelolechia pilati, 142+
Aerobryidium subpiligerum, 430+
Aerobryopsis subdivergens subsp. subdivergens, 128+, subsp. scariosa, 128+
Aerobryum speciosum, 128+
Agonimia tristicula, 56+
Agyrophora lyngei, 142+
AHMADJIAN, V. [Review:] G. G. Feige & H. T. Lumbsch (eds.) Phytochemistry and chemotaxonomy of lichenized ascomycetes—A festschrift in honour of Siegfried Huneck, 471
AHTI, T. *See* THOMSON, J. W. & T. AHTI, 138
Alectoria, 397+; nigricans, 44+, 142+; ochroleuca, 44+, 142+; sarmentosa, 142+, 399+; vancouverensis, 399+
Allantoparmelia almquistii, 142+; alpicola, 142+
ALLEN, B. [Review:] M. O. Hill et al. (eds.) Atlas of the bryophytes of Britain and Ireland, 349
ALLEN, B. *See* MAGILL, R. E. & B. ALLEN, 104, 338
ALLEN, B. *See* REDFEARN, P. L., JR., B. ALLEN & S. HE, 275
ALLEN, B. & R. E. MAGILL. Recent literature on mosses—160, 208;—162, 451
Aloina aloides, 415
Amygdalaria elegantior, 142+; panaeola, 142+; pelobotryon, 142+; subdissentiens, 142+
Anaptychia bryorum, 142+; fusca, 142+; hypoleuca, 409+; kaspica, 56+; palmulata, 409+, 426+; setifera, 56+
Anastrophyllum minutum, 168+
ANDERSON, L. E. [Review:] A. J. Sharp et al. (eds.) The moss flora of Mexico, 462
ANDERSON, L. E. *See* CRUM, H. & L. E. ANDERSON, 280
ANDREAS, B. K. [Review:] R. L. Stuckey. Women botanists of Ohio born before 1900, 218
Andromeda polifolia, 158, 284
Aneura, 422+, subg. Austroaneura, 64+; blasioides, 87+; **marianensis** sp. nov., 87*; maxima, 87+
Anoectangium thomsonii, 128+
Anomobryum filiforme var. concinnatum, 128+
Anomodon acutifolius, *275*; giraldii, 128+; minor subsp. integerrimus, 128+; thraustus, 128+
Antitrichia, 398+; curtipendula, 399+
Anzia ornata, 409+
Apatoplaca, 184+
APTROOT, A. [Review:] P. M. McCarthy. Saxicolous species of Porina Müll. Arg. (Trichotheliaceae) in the Southern Hemisphere, 467
Aptychella planula, *130*; yunnanensis, *130*
Arachniopsis coactilis, 71+; diacantha, 71+*; minima, 71+*
Arbutus menziesii, 40+
Arctoparmelia centrifuga, 44+, 142+; incurva, 44+, 142+; separata, 44+, 142+; subcentrifuga, 44+
Arthonia byssacea, 426+; didyma, 426+
Arthothelium ruanum, 426+
Arthrorhaphis alpina, 142+; citrinella, 142+
ARUP, U. The genus Caloplaca on seashore rocks in eastern North America, 377
Asahinea chrysantha, 44+, 142+; scholanderi, 44+, 142+
Aspicilia, 183+; aliena, 142+; anseris, 142+; aquatica, 142+; caesiocinerea, 142+; caesiopruinosa, 142+; cinerea, 142+; cf. cingulata, 142+; disserpens, 142+; fimbriata, 143+; heteroplaca, 143+; karelica, 143+; melanaspis, 143+; myrinii, 143+; plicigera, 143+; quartzitica, 143+; reptans, 56+; ryrkaipiae, 143+; subplicigera, 143+; supertegens, 143+
Astronia, 80
Athrotaxis selaginoides, 301
Atrichum crispulum, 128+; longifolium, 128; rhystophyllum, 128+
Aulacomnium, 135+; androgynum, 227+; heterostichum, 128+, 227+; palustre, 227+, 357+

Bacidia albidoplumbea, 298+; bagliettoana, 143+; beckhausii, 143+; herbarum, 143+; rubella, 426+
Baeomyces carneus, 143+; placophyllus, 143+; roseus, 143+; rufus, 143+
Banksia marginata, 298+
Barbilophozia binsteadii, 170+; floerkei, 170+; kunzeana, 167+; lycopodioides, 170+; quadriloba, 170+
Barbula fuscinervi, 257+; subcomosa, 129+; unguiculata, 129+
BARTHOLOMEW-BEGAN, S. E. & B. J. CRANDALL-STOTLER. The sporeling ontogeny of Monoclea gottschei subsp. elongata, 244
Bartramia halleriana, 129+; pomiformis, 129+, 227+
Bartramidula bartramioides, *133*; griffithiana, *133*
Bazzania adnexa var. **aucklandica** comb. & stat. nov., 319; convexa, *317*; cunninghamii, *319*; hochstetteri, 317+; involuta var. involuta, 314+, var. **submutica** comb. & stat. nov., 314; monilinervis, 314+*; nidita, 313+; **nova** sp. nov., 313*; taylorii, 313+; trilobata, 22
Bellemerea alpina, 143+; cinereorufescens, 143+
Biatora byssacea, 296+; carneoalbida, 143+; vernalis, 143+
Biatoropsis usnearum, 56+

[1] New scientific names and combinations are in boldface; page numbers of synonyms are in italic; taxa mentioned several times in an article are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

0007-2745/94/476–486$1.25/0

The Bryologist 97(4), 1994, pp. 476–486
Copyright © 1994 by the American Bryological and Lichenological Society, Inc.

Index to Volume 97 of THE BRYOLOGIST[1]

PIERS MAJESTYK
Biology Department, University of Southwestern Louisiana, Lafayette, LA 70504-2451

Abrothallus parmeliarum, 156+
Acacia mearnsii, 298+
Acarospora chlorophana, 142+; fuscata, 142+; glaucocarpa, 142+; gwynni, 440+; schleicheri, 142+; smaragdula, 142+; superfusa, 142+; veronensis, 142+
Acrocryphaea, 89+
Acromastigum colensoana, 319+; cunninghamii, 319+*; mooreanum, 314
Acroscyphus sphaerophoroides, 56+
Actinogyra muehlenbergii, 142+
Actinothuidium hookeri, 128+
Adelolechia pilati, 142+
Aerobryidium subpiligerum, 430+
Aerobryopsis subdivergens subsp. subdivergens, 128+, subsp. scariosa, 128+
Aerobryum speciosum, 128+
Agonimia tristicula, 56+
Agyrophora lyngei, 142+
AHMADJIAN, V. [Review:] G. G. Feige & H. T. Lumbsch (eds.) Phytochemistry and chemotaxonomy of lichenized ascomycetes—A festschrift in honour of Siegfried Huneck, 471
AHTI, T. *See* THOMSON, J. W. & T. AHTI, 138
Alectoria, 397+; nigricans, 44+, 142+; ochroleuca, 44+, 142+; sarmentosa, 142+, 399+; vancouverensis, 399+
Allantoparmelia almquistii, 142+; alpicola, 142+
ALLEN, B. [Review:] M. O. Hill et al. (eds.) Atlas of the bryophytes of Britain and Ireland, 349
ALLEN, B. *See* MAGILL, R. E. & B. ALLEN, 104, 338
ALLEN, B. *See* REDFEARN, P. L., JR., B. ALLEN & S. HE, 275
ALLEN, B. & R. E. MAGILL. Recent literature on mosses—160, 208;—162, 451
Aloina aloides, 415
Amygdalaria elegantior, 142+; panaeola, 142+; pelobotryon, 142+; subdissentiens, 142+
Anaptychia bryorum, 142+; fusca, 142+; hypoleuca, 409+; kaspica, 56+; palmulata, 409+, 426+; setifera, 56+
Anastrophyllum minutum, 168+
ANDERSON, L. E. [Review:] A. J. Sharp et al. (eds.) The moss flora of Mexico, 462
ANDERSON, L. E. *See* CRUM, H. & L. E. ANDERSON, 280
ANDREAS, B. K. [Review:] R. L. Stuckey. Women botanists of Ohio born before 1900, 218
Andromeda polifolia, 158, 284
Aneura, 422+, subg. Austroaneura, 64+; blasioides, 87+; **marianensis** sp. nov., 87*; maxima, 87+
Anoectangium thomsonii, 128+
Anomobryum filiforme var. concinnatum, 128+
Anomodon acutifolius, *275*; giraldii, 128+; minor subsp. integerrimus, 128+; thraustus, 128+
Antitrichia, 398+; curtipendula, 399+
Anzia ornata, 409+
Apatoplaca, 184+
APTROOT, A. [Review:] P. M. McCarthy. Saxicolous species of Porina Müll. Arg. (Trichotheliaceae) in the Southern Hemisphere, 467
Aptychella planula, *130*; yunnanensis, *130*
Arachniopsis coactilis, 71+; diacantha, 71+*; minima, 71+*
Arbutus menziesii, 40+
Arctoparmelia centrifuga, 44+, 142+; incurva, 44+, 142+; separata, 44+, 142+; subcentrifuga, 44+
Arthonia byssacea, 426+; didyma, 426+
Arthothelium ruanum, 426+
Arthrorhaphis alpina, 142+; citrinella, 142+
ARUP, U. The genus Caloplaca on seashore rocks in eastern North America, 377
Asahinea chrysantha, 44+, 142+; scholanderi, 44+, 142+
Aspicilia, 183+; aliena, 142+; anseris, 142+; aquatica, 142+; caesiocinerea, 142+; caesiopruinosa, 142+; cinerea, 142+; cf. cingulata, 142+; disserpens, 142+; fimbriata, 143+; heteroplaca, 143+; karelica, 143+; melanaspis, 143+; myrinii, 143+; plicigera, 143+; quartzitica, 143+; reptans, 56+; ryrkaipiae, 143+; subplicigera, 143+; supertegens, 143+
Astronia, 80
Athrotaxis selaginoides, 301
Atrichum crispulum, 128+; longifolium, 128; rhystophyllum, 128+
Aulacomnium, 135+; androgynum, 227+; heterostichum, 128+, 227+; palustre, 227+, 357+

Bacidia albidoplumbea, 298+; bagliettoana, 143+; beckhausii, 143+; herbarum, 143+; rubella, 426+
Baeomyces carneus, 143+; placophyllus, 143+; roseus, 143+; rufus, 143+
Banksia marginata, 298+
Barbilophozia binsteadii, 170+; floerkei, 170+; kunzeana, 167+; lycopodioides, 170+; quadriloba, 170+
Barbula fuscinervi, 257+; subcomosa, 129+; unguiculata, 129+
BARTHOLOMEW-BEGAN, S. E. & B. J. CRANDALL-STOTLER. The sporeling ontogeny of Monoclea gottschei subsp. elongata, 244
Bartramia halleriana, 129+; pomiformis, 129+, 227+
Bartramidula bartramioides, *133*; griffithiana, *133*
Bazzania adnexa var. **aucklandica** comb. & stat. nov., 319; convexa, *317*; cunninghamii, *319*; hochstetteri, 317+; involuta var. involuta, 314+, var. **submutica** comb. & stat. nov., 314; monilinervis, 314+*; nidita, 313+; **nova** sp. nov., 313*; taylorii, 313+; trilobata, 22
Bellemerea alpina, 143+; cinereorufescens, 143+
Biatora byssacea, 296+; carneoalbida, 143+; vernalis, 143+
Biatoropsis usnearum, 56+

[1] New scientific names and combinations are in boldface; page numbers of synonyms are in italic; taxa mentioned several times in an article are indicated by a plus sign (+) after the page number; illustrations are indicated by an asterisk (*).

0007-2745/94/476–486$1.25/0

Blasia, 64+
Blepharostoma trichophyllum, 167+
BONORA, A. *See* GERDOL, R., A. BONORA & F. POLI, 158
Botrydina viridis, 143+
BOWERS, F. D. [Review:] M. O. Hill et al. (eds.) Atlas of bryophytes of Britain and Ireland, 218
Brachysteleum microcarpum, *34*; polyphylloides, *37*
Brachythecium plumosum, 227+; rivulare, 129+
BRAKO, L. [Review:] R. Lücking. Foliicolous lichens—A contribution to the knowledge of the lichen flora of Costa Rica, Central America, 110
BREUSS, O. & B. MCCUNE. Additions to the pyrenolichen flora of North America, 365
Breutelia arundinifolia, 130+; deflexa, *130*; dicranacea, 129+; grandis, 421+; setschwanica, *129*; subdeflexa, *130*; wainioi, 421+; yunnanensis, *129*
Brigantiaea fuscolutea, 143+
BRODO, I. M. [Review:] G. Rambold & D. Triebel. The inter-lecanoralean associations, 111
BRODO, I. M. [Review:] R. Santesson. The lichens and lichenicolous fungi of Sweden and Norway, 468
Brodoa oroarctica, 143+
Brotherella recurvans, 22+
Bryocaulon divergens, 44+, 143+
Bryohaplocladium, *174*; angustifolium, *174*
Bryonora castanea, 143+; curvescens, 143+
Bryonorrisia acutifolia, 275+
Bryopteris diffusa, 422+
Bryoria, 397+; capillaris, 143+, 399+, 426+; chalybeiformis, 44+, 143+; fremontii, 143+; friabilis, 143+, 399+; fuscescens, 44+, 143+; glabra, 143+; lanestris, 44+, 143+; nadvornikiana, 44+; nitidula, 44+, 143+; oregana, 399+; pseudofuscescens, 143+, 399+; simplicior, 44+, 143+; tenuis, 44+; trichodes subsp. americana, 143+
Bryoxiphium norvegicum, 130+
Bryum argenteum, 130+; bicolor, 415; cavum, 421+; corrugatum, 421+, var. niveum, 421+; horizontale, 421+; pseudotriquetrum, 14+, 227+
BUCK, W. R. A nomenclatural correction in the Meteoriaceae, 436
BUCK, W. R. *See* THIERS, B. M. & W. R. BUCK, 91, 326
BUCK, W. R. The resurrection of Orthostichella, 434
BUCK, W. R. Three additions to the United States moss flora, 89
BUCK, W. R. Vainio's Brazilian bryophytes at TUR, 420
BUCK, W. R. & B. M. THIERS. Recent literature on hepatics and anthocerotes—136, 193;—138, 447
Buellia, 183+; coniops, 143+; disciformis, 143+, 298+; elegans, 143+; papillata, 143+; punctata, 143+; schaereri, 426+
BURK, W. R. & R. A. STUCKEY. Appendix. Publications and reports of Emanuel D. Rudolph (arranged chronologically), 443

CAI, HUAN-ZHONG. *See* ZHU, RUI-LIANG, LI-XIN YE & HUAN-ZHONG CAI, 277
Calamagrostis villosa, 284
Caliciales, 424
Calicium abietinum, 143+; salicinum, 301+, 426+; trabinellum, 426+; victorianum, 301+
Callicladium haldanianum, 22
Calluna vulgaris, 158
Calophyllum, 80
Caloplaca, 182+; ammiospila, 144+; approximata, 382+; arizonica, 439+; atrosanguinea, 144+; aurea, 184+; borealis, 144+; brattiae, 388+; brouardii, 184+; castellana, 382+; cerina, 144+; cinnamomea, *144*; cladodes, 185+; crenularia, 144+; discolor, 144+; ferruginea, *380*; flavovirescens, 382+; fraudens, 144+, 377+*; gloriae, 391+; granulosa, 391+; holocarpa, 144+; invadens, 144+; jungermanniae, 144+; lithophila, 377+*, var. elaeophora, *382*; luteominia, 382+; marina, 377+; microthallina, 377+*; nivalis, 144+; paulii, 184+; pyracea, 144+, 383+; scopularis, 377+*; sinapisperma, 144+; siphonospora, 184; stillicidiorum, 144+; tetraspora, 144+; thallincola, 391+; tucumanensis, 185+; verruculifera, 184+, 377+*; vitellinula, 383+; **wetmorei** sp. nov., 182*
Calymperes clemensiae, *80*; crassinerve, 376; erosum, 325; lonchophyllum, 80+; motleyi, 376; palisotii, 325; pallidum, 89+; serratum, 80+; subdisciforme, 371+, *416*; subserratum, 80+*; subulatum, 80+
Calymperopsis crenulata, 417+; martinicensis, *375*; subdisciformis, 376
Calypogeia, 252+; azurea, 167+
Calyptrochaeta japonica, 275+
Campanula rotundifolia, 83
Campylodontium regnellianum, 421+
Campylopodiella himalayana, 275+
Campylopodium khasianus, 132+
Campylopus arctocarpus, 421+; atrovirens, 130+; ericoides, 275+; flexuosus, 130+, 275+; fragilis subsp. goughii, 275+; laxitextus, 275+; penicillatus, 421+; polytrichoides, 421+; schmidii, 275+; strictifolius, 421+; subarctocarpus, 421+
Candelaria concolor, 144+; fibrosa, 409+
Candelariella athallina, 144+; aurella, 144+; dispersa, 144+; lutella, 144+; placodizans, 144+; terrigena, 144+; vitellina, 144+; xanthostigma, 144+; xanthostigmoides, 298+
Candida albicans, 124
CANO, M. J., J. GUERRA & R. M. ROS. Pterygoneurum compactum sp. nov. (Musci: Pottiaceae) from Spain, 412
CAO, TONG & D. H. VITT. North American–East Asian similarities in the genus Ptychomitrium (Bryopsida), 34
Carbonea atronivea, 144+; vorticosa, 144+
CÁRDENAS S., A. & C. DELGADILLO M. Lorentziella imbricata and Oreoweisia delgadilloi in Mexico, 85
Carex, 83; curta, 285; limosa, 284; pauciflora, 158; rostrata, 284+
Castanopsis acuminatissima, 80
Catapyrenium acarosporoides, 366+; andicola, 365+; chilense, 365+; cinereum, 57+, 144+; compactum, 144+, 366+; **congestum** sp. nov., 365+; contumescens, 366+; daedaleum, 56+, 366+; lachneum, 144+, 365+; lacinulatum, 367+; norvegicum, 365+; podolepis, 365+; psoromoides, 365+; rufescens, 365+; squamulosum, 57+, 367+; tuckermanii, 366+; umbrinum, 365+; waltheri, 365+
Catillaria, 298+; glauconigrans, 144+; rudolphi, 440+
Catinaria atropurpurea, 426+; laureri, 426+
Cavernularia hultenii, 144+; lophyrea, 144+
"Celidium" insitivum, 156+
Cephalozia bicuspidata, 167+; lunulifolia, 168+; ovalifolia, 422+; pleniceps, 169+
Cephaloziella exiliflora, 14+
Ceratodon purpureus, 14+
Cetraria andrejevii, 44+; californica, 144+; chlorophylla, 399+; commixta, 44+, 144+; cucullata, 44+, 83+, 144+; delisei, 44+, 144+; ericetorum, 83+, 144+, var. reticulata, 45+; fastigiata, 45+, 144+; hepatizon, 45+, 144+; islandica, 45+, 83+, subsp. crispiformis, 144+, subsp. islandica, 144+; kamczatica, 144+; laevigata, 45+, 144+; nigricans, 45+, 144+; nigricascens, 45+, 144+; nivalis 45+, 83+, 144+; orbata,

399+; pallidula, 399+; platyphylla, 399+; sepincola, 45+; subalpina, 144+; tilesii, *83*
Cetrelia alaskana, 144+; chicitae, 426+
Chaenotheca brunneola, 426+; cinerea, 57+; furfuracea, 426+; trichialis, 426+; xyloxena, 145+
Chaenothecopsis debilis, 426+; pusilla, 426+
Chamaebryum, 85+
CHAPMAN, R. L. [Review:] L. E. Graham. Origin of land plants, 346
Cheilolejeunea imbricata, 277+
Chrysocladium flammeum, 130+; phaeum, 130+
Citrus aurantium, 405+
Cladina, 69+; arbuscula, 45+, 145+; mitis, 45+, 145+; rangiferina, 45+, 145+; stellaris, 45+, var. aberrans, 145+, var. stellaris, 145+; stygia, 45+, 145+
Cladonia, 67+, sect. Cladonia, 67+, sect. Cocciferae, 69+, sect. Perviae, 67+; acuminata, 45+, 145+; alaskana, 42+, 145+; amaurocraea, 45+, 145+; artuata, 68+; bacillaris, 58+; bacilliformis, 145+; bellidiflora, 45+, 145+; borealis, 145+; botrytes, 145+; cariosa, 45+, 145+; carneola, 45+, 145+; cenotea, 45+, 145+; cervicornis, 67+, subsp. verticillata, 68+; chlorophaea, 45+, 68+, 117+, 145+; clavulifera, 409+; coccifera, 45+; coniocraea, 145+; cornuta, 45+, subsp. cornuta, 145+, subsp. groenlandica, 145+; crispata, 145+, var. cetrariiformis, 145+; cristatella, 118+; cryptochlorophaea, 26+, 42+; cyanipes, 45+, 145+; decorticata, 145+; deformis, 45+, 145+; digitata, 45+, 145+; dimorpha, 68+; ecmocyna, 45+, 68+, 145+, subsp. intermedia, 145+; fimbriata, 45+, 68+, 145+, 399+; firma, 68+; furcata, 68+, 145+; gracilis, 45+, subsp. gracilis, 145+, subsp. elongata, 145+, subsp. turbinata, 145+, subsp. vulnerata, 145+; grayi, 45+, 68+, 145+; kanewskii, 145+; luteoalba, 56+, 145+; macilenta, 68+, 145+; macroceras, 145+; macrophylla, 45+, 145+; macrophyllodes, 45+, 145+; maxima, 45+, 146+; merochlorophaea, 45+, 68+, 146+, 399+, var. merochlorophaea, 69+; metacorallifera, 45+, 146+; multiformis, 68+, 146+; nipponica, 146+; ochrochlora, 68+, 146+, 399+; phyllophora, 146+; pleurota, 45+, 146+; pocillum, 45+, 146+; prolifica, 68+; pseudorangiformis, 146+; pseudostellata, 146+; pyxidata, 45+, 146+; rei, 146+; scabriuscula, 45+, 68+, 146+; squamosa, 45+, 146+, var. subsquamosa, 68+, 399+; stricta, 45+, 146+; subcervicornis, 45+; subfurcata, 45+, 146+; submitis, 409+; subulata, 146+; sulphurina, 45+, 146+; symphycarpa, 45+, 146+; thiersii, 68+; thomsonii, 146+; transcendens, 68+; turgida, 146+; uncialis, 45+, 146+; verruculosa, 68+, 146+; verticillata, 45+, 146+; wainii, 146+
Clastobryopsis planula, 130+
Clematis fremontii, 124
Climacium dendroides, 130+
Cliostomum griffithii, 146+
Coccotrema cucurbitula, 298+
Cochliobolus heterostrophus, 124
Coelocaulon aculeatum, 45+, 146+; muricatum, 146+
Coenogonium, 186+; boninense, 186+; deightoni, 186+; leprieurii, 186+; **missouriense** sp. nov., 186•; moniliforme, 186+
Collema auriforme, 56+; bachmanianum, 146+; crispum, 45+, 146+; curtisporum, 146+; furfuraceum, 146+; glebulentum, 45+, 146+; nigrescens, 426+; polycarpon, 146+; subflaccidum, 146+, 426+; tenax, 45+, 146+; undulatum var. granulosum, 146+
Collemopsidium pyrenuloides, 440+
Cololejeunea appressa, 277+; goebelii, 277+; inflata, 277+; latilobula, 278+; latistyla, 278+; longifolia, 278+; longilobula, 278+; macounii, 278+; ocelloides, 278+; ornata, 278+; oshimensis, 277+; planissima, 278+; pseudofloccosa, 278+; raduliloba, 278+; spinosa, 278+; subkodame, 278+; subocelloides, 278+; verdoornii, 278+
Colura tenuicornis, 278+
Conocephalum, 9+
Conomitrium, 253+; amoenum, *254*; asterodontium, *262*; brunneolum, *254*; caripense, *266*; elachistophyllum, *260*; excavatum, *258*; falcatus, *255*; glaziovii, *259*; goebelii, 255+; hemiloma, *266*, var. atrovirens, *258*; intramarginatum, *260*; longipedicellatum, 421+; perpygmaeum, *260*; platybryoides, *254*; platyphyllus, *254*; polycarpum, *259*; psatyrocheilon, *265*; puiggarii, *264*; serratus, 254+; smaragdinum, *255*; trachelymum, *264*; variabilis, *255*
COOPER-ELLIS, S. [Review:] E. C. Pielou. The interpretation of ecological data, 468
Coprinus cinereus, 118+
Costesia, 85+
Cotula maniototo, 65+
CRANDALL-STOTLER, B. [Review:] A. K. Asthana & S. C. Srivastava. Indian hornworts (a taxonomic study), 112
CRANDALL-STOTLER, B. J. *See* BARTHOLOMEW-BEGAN, S. E. & B. J. CRANDALL-STOTLER, 244
Crassicosta, *174*; hermanii, *174*
Cratoneuron filicinum, 130+
CROSBY, M. R. *See* TAN, B. C., QI-WEI LIN, M. R. CROSBY & PAN-CHENG WU, 127
Crossidium chloronotis, 275+; crassinerve, 415
CRUM, H. & L. E. ANDERSON. A deviant expression of Sphagnum tenerum from the Virginia tidewater and the Dominican Republic, 280
Cryphaea, 89+; filiformis, 89+
Cryptopodium bartramioides, 227+
Curvicladium, 305+
Cyathea, 314
Cyathodes parvifolia, 298+
Cyathophorella hookeriana, 130+; intermedium, *130*; tonkinensis, 130+
Cyclodictyon laetevirens, 227+
Cyphelium tigillare, 146+
Cystocoleus, 186+; niger, 186+
CZECZUGA, B. & S. HAMMER. Carotenoids in representatives of the lichen genus Cladonia in North America, 67

Dactylina arctica, 45+, 146+; beringica, 45+, 146+; madreporiformis, 45+, 146+; ramulosa, 45+, 146
Dactylospora lobariella, 56+
Daphnia pulex, 122+
DAVIS, J. S. Coenogonium missouriense, a new lichen species from Missouri, 186
Decodon brasiliensis, 421+
Degelia plumbea, 146+
DELGADILLO M., C. *See* CÁRDENAS S., A. & C. DELGADILLO M., 85
Dendriscocaulon intricatulum, 146+
Dendrocyathophorum herzogii, 130+; paradoxum, *130*
DEPRIEST, P. T. [Review:] D. J. Galloway. (ed.) Tropical lichens: their systematics, conservation, and ecology, 347
DEPRIEST, P. T. Variation in the Cladonia chlorophaea complex II: Ribosomal DNA variation in a southern Appalachian population, 117
Dermatocarpella, 367+
Dermatocarpon intestiniforme, 146+; lorenzianum, 365+; miniatum, 146+; rivulorum, 45+
Dicranella exigua, 421+; fusca, 421+; martii, 421+; nitida, 421+

Dicranoloma cylindrothecium, 128+
Dicranum flagellare, 20+, 358+; fuscescens, 17+, 399+; scoparium, 22+
Didymodon fallax, 130+; giganteus, 130+
DIEDERICH, P. *See* GOWARD, P., P. DIEDERICH & R. ROSENTRETER, 56
Dimelaena oreina, 146+
Dimerella pineti, 426+
Diphyscium fulvifolium, 130+; involutum, 130+
Diplophyllum albicans, 167+; taxifolium, 168+
Diploschistes muscorum, 146+; scruposus, 147+
Diselma archeri, 301
Distichophyllum carinatum, 275+
Ditrichum crispatissimum, 130+; flexicaule, 130+; heteromallum, 130+; pallidum, 130+; subrufescens, 421+
Dolichomitra cymbifolia, 130+
Douglasia laevigata, 83
Douinia ovata, 169+
Drepanocladus aduncus, 357+; fluitans, 284
Drepanolejeunea angustifolia, 278+; erecta, 278+
Drosera, 280
Duthiella declinata, 227+; flaccida, 130+; speciosissima, 227+; wallichii, 227+

Ectropothecium subdenticulatum, 421+; wainioi, 421+
Edrudia, 442+
Elaeocarpus, 80
ELIX, J. A. *See* KANTVILAS, G. & J. A. ELIX, 296
Endocarpon adscendens, 368+; adsurgens, 365+; pallidum, 57+; pusillum, 368+; tortuosum, 45+
ENGEL, J. J. & G. L. SMITH-MERRILL. Studies of New Zealand Hepaticae. 8–13. Bazzania and Acromastigum, 313
ENGEL, J. J. & R. M. SCHUSTER. Studies of New Zealand Hepaticae. 7. The status of Steereomitrium, 63
ENROTH, J. *See* HYVÖNEN, J. & J. ENROTH, 305
ENROTH, J. & R. E. MAGILL. Neckeropsis pocsii (Neckeraceae, Musci) a new species from Comoro Islands, 171
Entodon conchophyllus, *134*
Entosthodon buseanus, 130; wichurae, 130
Ephebe hispidula, 45+, 147+; lanata, 45+, 147+
Ephelidium, 440+; heardense, 440+
Epilichen scabrosus, 147+
Eriophorum angustifolium, 284+; vaginatum, 158, 284
Erpodium biseriatum, 128+
ESSLINGER, T. L. Recent literature on lichens—153, 94;—154, 196;—155, 330
Esslingeriana idahoensis, 399+
Eumyurium sinicum, 131+
Euopsis granatina, 147+
Eurhynchium oreganum, 399+; pulchellum, 358+
Evernia esorediosa, 176+; mesomorpha, 45+, 147+; perfragilis, 45+, 147+; prunastri, 399+

Fabronia ciliaris, 128+
Fauriella tenerrima, 131+; tenuis, 131+
Fissidens sect. Bryoidium, 266+, sect. Crenularia, 258+, sect. Pachylomidium, 255+, sect. Pycnothallia, 266+, sect. Semilimbidium, 258; subg. Eufissidens, 253+, subg. Fissidens, *253*; acicularis, *260*; acutissimus, *267*; adelphinus, 131+; allenianus, *264*; allionii, 254+; alticaulis, *256*; altolimbatus, *265*; amoenus, 253+; andicola, 254; anguste-limbatus, 255+; angustelimbatus var. brevisetus, *255*; angustifolius, 255+; anomalus, 131+; antennidens, *264*; areolatus, *131*; asplenioides, 258+; asterodontius, *262*; austro-adiantoides, *255*; balansaeanus, *266*; bockii, *257*; bourgaeanus, 255+; brevifalcatus, *260*; brevipes, 256+; brotheri, *257*; brotherianus, *257*, var. longisetus, *257*; brunneolus, *254*; bryoides, 253+, var. bryoides, 131+, 253+, var. lateralis, 131+; buckii, *264*; caespitulosus, *259*; camptothecius, *260*; capillisetoides, *255*; capillisetus, *257*; caripensis, *266*; catharinensis, *254*; catumbensis, *261*; celatocognatus, *258*; ceylonensis, 131+; chilensis, *256*; chrysopoma, *256*; circinatulus, *256*; constrictus, *260*; crassicuspis, *257*; crassipes, 131+; crenatulus, *261*; crispo-falcatulus, *258*; crispus, 253+; cristatus, *131*; curvatus, 256+; cuspidatus, 267+; cuspidulatus, *258*; densiretis var. latifolius, *258*; diplodus var. wainionis, 257+; distichellus, *264*+; diversiformis, *257*; divisus, *262*; dubius, 131+, 255+; duryae, *255*; elachistophyllus, *268*; elegans, 253+; evanescens, *259*; excavatus, *258*; excurrentinervis, 257+; falcatus, *255*; faucium, *265*; felipponei, *264*; fendleri, *256*; fernandezianus, *256*; filirameus, *257*; flaveolus, 256+; flavifrons, *260*; flavinervis, *258*; flavosetus, *257*; flexinervis, *262*; francii, *257*; fratis, *258*; garberi, *258*; geheebii, 259+; geminidens, *258*; geppii, 128+; glaucifrons, *258*; glaziovii, *259*; goebelii, *255*; goyazensis, 259+; grandifrons, 131+; gymnogynous, 131+; guarani, *259*; guarapiensis, *266*; guianensis, 258+; hampeanus, *266*; hauthalii, *264*, var. angustissimus, *264*, var. gracillima, *267*, var. minor, *264*, var. obtusatula, *264*; hemibryoides, *260*; hemicraspedophyllus, *266*; hemiloma, *266*, var. atrovirens, *258*; hornschuchii, 259+; hyalinus, 131+; inaequalis, 260+; incacorralis, *260*; incertus, 265+; incisus, 257+; indistinctus, *257*, var. major, *267*; innovans, *266*; intramarginatus, 260+; involutus, 131+; jujuiensis, *267*; kochii, *265*; latifolius, 262, *262*; laxe-reticulatus, *257*; laxus, *262*; leptocaulis, *255*; leptophyllus, 261+, *261*; leptopodus, *258*; leucopteris, *266*; lindigii, *257*; linearis, 259+; lonchothecius, *256*; longicaulis, *266*; longipedicellatus, *259*; luteo-limbatus, *266*; luteo-viridis, *256*; macroblastus, *260*; mangarevensis, *131*; mararyensis, *267*; micropyxis, *261*; minutissimus, *257*; minutus, 258+; mollis, 261+; monomorphus, *258*; nanoloma, *258*; neglectus, 259+; neonii, *264*; nicholsonii, *266*; nobilis, 131+; oblongifolius, 131+; obscurirete, 131+; obscurus, 131+; obtusifolius, 258+; occultus, *260*; oediloma, 261+, var. steyermarkii, *263*; oerstedianus, *262*+; oligophyllus, *256*; opacus, *266*; osmundioides, 258+; ovicarpus, *259*; pabstii, *258*, var. latifolius, *261*; papilliferus, *262*; papillosus, *258*; paraguensis, *259*; paulensis, *258*; pauperculus var. pauperculus, *260*, var. surinamensis, *260*; pellucidus var. **asterodontius** comb. nov., 253, var. **papilliferus** comb. nov., 253, var. pellucidus, 261+; pennula, *255*; percrispus, *256*; perdecurrens, 131+; perexilis, *257*; perfalcatus, 262+; perminutus, *260*; perpygmaeus, *260*; plagiochiloides, 131+; platyphyllus, *254*; polycarpus, *259*; polypodioides, 131+, 253+; prionodes, 254+, fo. flexinervis, *262*, fo. hornschuchii, *262*, fo. puiggarii, *264*; prosenchymaticus, *255*; protracticaulis, *255*; psatyrocheilon, *265*; pseudoincurvatus, *256*, var. latifolius, *256*; pseudo-monomorphus, *260*; pseudorepandus, *266*; puiggarii, *264*; radicans, 263+; ravenelii, *258*; reclinatulus, *256*; reesei, *261*; repandus, *256*; rigidulus, 263+, subsp. masatierrensis, *263*; rubentiloma, *260*; rubiginosulus, *264*; saltograndensis, *260*; santa-clarensis, 264+, var. obtusifolius, *263*; saxicola, *266*; scalaris, 264+; scariosus, 253+; sehnemii, *264*; serratus, 264+; sharpii, *262*; similis, *266*; smaragdinus, *255*; squamulatus, *265*; steyermarkii, *263*; subaloma, *256*; subcrispus, *256*; subflexinervis, *261*; subinclinatulus, *257*; submarginatus, 265+; submicropyxis, *265*; subradicans, 265+; subulatus, 265+; taxifolius, 131+; tejoensis, *258*; te-

ner, *264*; tenerrimus, 421+; tenuifolius, *260*; teysmannianus, 131+; tocoraniensis, *264*; **townsendianus** nom. & stat. nov., 253; trachelyma, *264*; uncinatus, *261*; validicostatus, *265*; variabilis, *255*; wallisii, 265+; weirii var. hemicraspedophyllus, 266+, var. insertus, *265*, var. weirii, 266+; zippelianus, 131+; zollingeri, 266+
Fistulariella dilacerata, 147+; roesleri, 147+
Floribundaria setschwanica, 131+; sparsa, 131+
Fossombronia, 244+*
Franciella spiridentoides, 227+
Fraxinus angustifolia, 405+
Frullania beyrichiana, 422+; brasiliensis, 422+; flexicaulis, 422+; gibbosa, 422+; glomerata, 422+; hians, 422+; lindbergiana, 422+; moniliata, 278+; mucronata, 422+; muscicola, 278+; nisquallensis, 169+, 399+; sebastionopolitana, 422+; squarrosa, 422+
Fulgensia, 184+; bracteata, 147+; desertorum, 56+
Funaria buseana, *130*; hygrometrica, 131+
FURUKI, T. Aneura marianensis sp. nov. (Hepaticae) of the northern Mariana Islands, 87
Fuscidea, 298+

GARILLETI, R. *See* LARA, F., R. GARILLETI & V. MAZIMPAKA, 402
Geocalyx graveolens, 168+
GERDOL, R., A. BONORA & F. POLI. The vertical pattern of pigment concentrations in chloroplasts of Sphagnum capillifolium, 158
Gigaspermaceae, 85+
Gigaspermum, 85+; repens, 85+
GLEW, K. A. Range extension of the lichen Vulpicida tilesii (Ascomycotina, Parmeliaceae) into the Pacific Northwest of the United States, 83
Glyphomitrium angustifolium, *34*+; sinense, *34*+
GOFFINET, B. [Review:] J. Lewinsky. Monographic studies on Orthotrichum (Musci), 464
Gongylia nadvornikii, 147+
Goniobryum subbasilare, 227+
GOULD, W. A. Macrolichens of the Coppermine, Hood, and Thomsen rivers, Northwest Territories, Canada, 42
GOWARD, T., P. DIEDERICH & R. ROSENTRETER. Notes on the lichens and allied fungi of British Colombia. II, 56
Graphis scripta, 147+
GREVEN, H. C. Grimmia crassifolia new to Europe, 180
GRIFFIN, III, D. [Review:] N. Salazar Allen. A revision of the pantropical moss genus Leucophanes Brid., 217
Grimmia antarctica, 14+; crassifolia var. crassifolia, 180+*, var. cucullata, 180+; laevigata, 180+; obtusifolia, 275+; ovalis, 180+; tergestina, 180+
GUERRA, J. *See* CANO, M. J., J. GUERRA & R. M. ROS, 412
Gyalecta foveolaris, 147+
Gyalidea lecideopsis, 147+
Gymnomitrion coralloides, 168+
Gymnostomum aurantiacum, 131+

Haematomma infuscum, 298+; lapponicum, 147+; ochrophaeum, 409+
HAMADA, N., K. OKAZAKI & M. SHINOZAKI. Accumulation of monosaccharides in lichen mycobionts cultured under osmotic conditions, 176
HAMMER, S. *See* CZECZUGA, B. & S. HAMMER, 67
Haplocladium angustifolium, 174+; microphyllum, 131+; transvaaliense, 174+
Haplodontium, *1*+
Haplomitrium, 63, 244+*; hookeri var. hookeri, 63+, var. minutum, *65*; **minutum** comb. nov., 65*
HARADA, H. Trichothelium marianense, a new foliicolous lichen species from the Mariana Islands, Micronesia, 272
Harpanthus flotovianus, 167+
Hawksworthiana peltigericola, 56+
HE, S. *See* REDFEARN, P. L., JR., B. ALLEN & S. HE, 275
HEDENÄS, L. The basal pleurocarpous diplolepidous mosses—a cladistic approach, 225
Hedwigia ciliata, 131+
Helodium, 174+
Heppia lutosa, 58+
Herpetineuron toccoae, 131+
Heterodermia speciosa, 147+
Heteroscyphus, 314; coalitus, 278+
Hildebrandtiella cuspidans, 430+
Himantocladium warburgii, 132+
Himantormia, 17+
Holomitrium densifolium, 131+; griffithianum, *131*
Homalia lusitanica, 227+; subarcuata, *135*
Homaliadelphus targionianus, 131+
Homaliodendron exiguum, 127+; microdendron, 131+; neckeroides, 131+; scalpellifolium, 131+
Homalothecium nilgheriense, 430+
HONG, W. S. Tritomaria in western North America, 166
Hookeria acutifolia, 131+; merkelii, 421+; wainioi, 421+
Hookeriopsis geminidens, *131*; secunda, *131*; utacamundiana, 127+
HUHNDORF, S. M. [Review:] A. Aptroot. A monograph of the Pyrenulaceae (excluding Anthracothecium and Pyrenula) and the Requienellaceae, with notes on the Pleomassariaceae, the Trypetheliaceae and Mycomicrothelia (lichenized and non-lichenized Ascomycetes), 346
Hylocomium splendens, 127+
Hymenelia caracea, 395+; lacustris, 147+, 393+
Hymenodon pilifer, 227+
Hymenodontopsis stresemannii, 227+
Hyophila involuta, 132+
Hypnodendron dendroides, 227+; kerrii, 227+
Hypnum angustifolium, *174*; circinale, 399+; hexastichum, *435*; imponens, 22+; pentastichum, *435*
Hypocenomyce australis, 301+; castaneocinerea, 58+; friesii, 147+; scalaris, 147+
Hypogymnia, 397+; apinnata, 147+, 399+; austerodes, 45+, 147+; bitteri, 45+, 147+; enteromorpha, 147+, 399+; imshaugii, 147+, 399+; inactiva, 399+; lugubris, 301+; metaphysodes, 399+; occidentalis, 147+, 399+; oroarctica, 45+; physodes, 45+, 147+, 399+; subobscura, 45+, 147+; subphysodes, 301+; tasmanica, 298+; tubulosa, 147+, 399+, 426+; vittata, 147+
Hypopterygium arbuscula, 227+; didictyon, 227+; fauriei, *132*; japonicum, 132+
HYVÖNEN, J. & J. ENROTH. Cladistic analysis of the genus Pinnatella (Neckeraceae, Musci), 305

Icmadophila ericetorum, 147+
Ilex latifolia, 277+
Illosporium carneum, 58+
Imshaugia aleurites, 147+
India, Porina from, 73+
Indocalamus tessellatus, 277+
Ionaspis **alba** sp. nov., 393*; epulotica, 147+; odora, 393+
Isocladiella surcularis, 128+
Isopterygium pohliaecarpum, *134*; tenerifolium, 89+; textorii, *134*
Isotachis serrulata, 422+
Isothecium, 398+; myosuroides, 132+, 399+

Japewia tornoensis, 147+
Jungermannia adnexa, *317*; atrovirens, 167+; exsecta, *166*; monilinervis, *314*; scitula, *166*; sphaerocarpa, 169+

Kantia amazonica, 422+
KANTVILAS, G. & J. A. ELIX. Ramboldia, a new genus in the lichen family Lecanoraceae, 296
Kiliasia athallina, 147+
KIMMERER, R. W. Ecological consequences of sexual versus asexual reproduction in Dicranum flagellare and Tetraphis pellucida, 20

LARA, F., R. GARILLETI & V. MAZIMPAKA, Orthotrichum macrocephalum sp. nov., a new moss of section Diaphana from the Iberian Peninsula, 402
Lasallia pensylvanica, 45+, 147+
LAWREY, J. D. [Review:] C. D. Bonham. Measurements for terrestrial vegetation, 472
LAWREY, J. D. [Review:] P. W. Price et al. (eds.) Plant–animal interactions, 216
LAWREY, J. D. [Review:] D. W. Tallamy & M. J. Raupp. (eds.) Phytochemical induction by herbivores, 220
Lecanactis chloroconia, 426+
Lecania dubitans, 148+
Lecanora argopholis, 148+; atrosulphurea, 148+; beringii, 148+; boligera, 148+; cenisia, 148+; circumborealis, 148+; crenulata, 148+; elmorei, 439+; epibryon, 148+; expallens, 148+; fuscescens, 148+; gangaleoides, 148+; geophila, 148+; glabrata, 410; hagenii, 148+; hypoptoides, 148+; impudens, 148+; imshaugii, 409+; intricata, 148+; luteovernalis, 148+; marginata, 148+; megalocheila, 409+; microthallina, *384*; mughicola, 58+; nordenskioeldii, 148+; cf. pallida, 148+; perflexuosa, *409*; persimilis, 58+; polytropa, 148+; pulicaris, 148+; rupicola, 148+; salicicola, 148+; scopularis, *387*; straminea, 148+; strobilina, 148+; subimmergens, 409+; subintricata, 148+; subradiosa, 148+; symmicta, 148+; varia, 148+; zosteraе, 148+
Lecidea albohyalina, 148+; alpestris, 148+; aniptiza "*intersociella," *302*; atrobrunnea, 148+; auriculata, 148+; cinnabarina, 148+; confluens, 148+; cuprea, 148+; diapensiae, 148+; elabens, *296*; erythrophaea, 148+; fuscoatra, 148+; immarginata, 298+; lactea, 148+; lampra, *296*; lapicida fo. lapicida, 148+, fo. ochracea, 148+; leptoboloides, 148+; lithophila, 148+; lulensis, 148+; melaphanoides, 148+; myriocarpella, 148+; nylanderi, 426+; paupercula, 148+; petraeoides, *298*; picea, 148+; plebeja, 148+; plicatula, *299*; ramulosa, 148+; shushanii, 148+; stenotera, 148+; stuartii, *296*; cf. subcandida, 148+; subnexa, *296*; tessellata, 148+; umbonata, 148+
Lecidella carpathica, 148+; elaeochroma, 148+; euphorea, 148+, 426+; stigmatea, 148+; wulfenii, 148+
Lecidoma demissum, 149+
Leciophysma finmarkicum, 58+, 149+
Lejeunea anoplantha, 422+; brotheri, 422+; catanduana, 278+; contigua, 422+; devoluta, 423+; didrichseni, 423+; flava, 278+; muelleriana, 423+; parva, 278+; rionegrensis, 423+; saccatiloba, 423+; stenophylla, 423+; sterilis, 423+; ulicina, 278+; unciloba, 423+
Lepidozia brevifissa, 423+; cupressina, 422+; reptans, 170+; setigera, 314+
Lepraria caesioalba, 149+; lobificans, 149+
Leprocaulon, 83+; subalbicans, 149+
Leproloma cacuminum, 149+
Leptocolea longilobula, 278+
Leptodontium scaberrimum, 132+; viticulosoides, *132*
Leptogium sp., 45+; brebissonii, 56+; crenulatum, 45+; cyanescens, 149+; hirsutum, 149+; lichenoides, 149+; saturninum, 149+, 426+; schraderi, 56+; tenuissimum, 45+, 149+; teretiusculum, 149+
Leptolejeunea elliptica, 278+; yangii, 279+
Leptotheca gaudichaudii, 227+
Leskea imponderosa, *436*
Leucobryum candidum, 314+; sordidum, 421+; widgrenii, 421+
Leucodon sinense, 132+
Leucolepis menziesii, 227+
Leucoloma biplicatum, 421+; mittenii, 276+
LI, YENHUNG & D. H. VITT. The dynamics of moss establishment: temporal responses to nutrient gradients, 357
Libocedrus bidwillii, 301
Lichenochora thallina, 56+
Lichenodium sirosiphoideum, 149+
Lichenosticta alcicornaria, 58+
LIN, QI-WEI *See* TAN, B. C., QI-WEI LIN, M. R. CROSBY & PAN-CHENG WU, 127
Lobaria, 57+; hallii, 149+, 399+; kurokawae, 149+; linita, 45+, 149+; oregana, 149+, 321+, 397+; pseudopulmonaria, 149+; pulmonaria, 57+, 149+, 399+, 426+; quercizans, 409+, 426+; scrobiculata, 149+, 399+
Locusta migratoria, 122+
Lopadium disciforme, 149+; pezizoideum, 149+
Lophocolea decurrens, 423+
Lophozia, 166+; aequifolia, 423+; ascendens, 170+; guttulata, 167+; heterocolpos, 167+; incisa, 167+; sudetica, 168+; ventricosa, 170+
Lorentziella imbricata, 85+
Loxospora elatina, 426+; ochrophaea, *409*
Loxotis, 436
Lupinus lepidus, 83
LUTZONI, F. M. Ionaspis alba (Ascomycotina, Hymeneliaceae), a new lichen species from eastern North America, 393

Macromitrium argutum, 421+; nitidum, 421+
Macrothamnium macrocarpum, 132+
Madotheca brasiliensis, 423+
MAGILL, R. E. *See* ENROTH, J. & R. E. MAGILL, 171
MAGILL, R. E. *See* REESE, W. D., R. E. MAGILL & T. PÓCS, 430
MAGILL, R. E. & B. ALLEN. Recent literature on mosses—159, 104;—161, 338
MARAVOLO, N. C. *See* STANISLAUS, R. C. & N. C. MARAVOLO, 162
MARCANO, V. & A. MORALES MÉNDEZ. New species of Ramalina from Venezuela, 26
MARCANO, V. & A. MORALES MÉNDEZ. Ramalina africana in Venezuela, 190
Marchantia, 162+; chenopoda, 423+; polymorpha, 162+
Masonhalea richardsonii, 45+, 149+
Massalongia carnosa, 45+, 149+; microphylliza, 56+
Mastigobryum acuminatum, 423+; arcuatum, 423+; cunninghamii, *319*; martianum, 423+; monilinerve, *314*; nitens, *317*; novae-hollandiae fo. aucklandicum, *319*, fo. submuticum, *314*; phyllobolum, 423+; vincentianum, 423+
MATA GARCÍA, L. B. Rimelia bonplandii, a new lichen species (Ascomycotina: Parmeliaceae), 432
MAZIMPAKA, V. *See* LARA, F., R. GARILLETI & V. MAZIMPAKA, 402
MCCUNE, B. *See* BREUSS, O. & B. MCCUNE, 365
MCCUNE, B. Using epiphyte litter to estimate epiphyte biomass, 396

McQueen, C. B. [Review:] A. L. Heathwaite. (ed.) Mires. Process, exploitation, and conservation, 473
McQueen, C. B. [Review:] N. G. Miller. (ed.) Biology of Sphagnum, 469
Megalaria, 301+
Megaspora verrucosa, 149+
Melanelia albertana, 149+; disjuncta, 45+, 149+; elegantula, 149+; exasperatula, 45+, 149+; infumata, 45+, 149+; multispora, 149+; olivacea, 149+; olivaceoides, 45+, 149+; panniformis, 45+, 149+; septentrionalis, 45+, 149+; sorediata, 45+, 149+; stygia, 45+, 148+; subaurifera, 149+; subglabra, 298+; tominii, 45+, 149+; trabeculata, 149+
Melick, D. R. & R. D. Seppelt. Seasonal investigations of soluble carbohydrates and pigment levels in antarctic bryophytes, 13
Melosira roeseana, 186+
Menegazzia myriotrema, 298+; terebrata, 399+; weindorferi, 298+
Merrill, G. L. Smith. *See* Engel, J. J. & G. L. Smith Merrill, 313
Mesochaete undulata, 227+
Meteoriella soluta, 132+
Meteoriopsis reclinata, 132+
Meteorium ambiguum, 421+; atrovariegatum, 132+; horikawae, 132+; miquelianum, 132+, subsp. atrovariegatum, *132*; papillarioides, 132+; piliferum, *132*; remotifolium, 421+; subpolytrichum, 132+; taiwanense, 132+; "widgrenii prox. remotifolium," 421+
Metzgeria conjugata, 423+; furcata, 279+
Micarea assimilata, 149+; crassipes, 149+; denigrata, 149+; globulosella, 303+; incrassata, 150+; **intersociella** comb. & stat. nov., 296; melaena, 150+, 426+; peliocarpa, 150+; prasina, 150+, 426+; synotheoides, 304+
Microcampylopus laevigatus, 128+
Microthamnium campaniforme, 421+; flavidum, 421+; widgrenii, 421+
Mielichhoferia, 1+; elongata, 1+, 47+; macrocarpa, 2+, 47+; mielichhoferiana, 1+, 47+; nitida, 2; tehamensis, 2, 47+
Miriquidica, 297+
Mitthyridium flavum, 430+; micro-undulatum subsp. **comoroensis** subsp. nov., 430*, subsp. micro-undulatum, 430+*
Miyawaki, H. Lecanora imshaugii, a lichen of eastern North America and eastern Asia, 409
Mnium hornum, 227+; laevinerve, 132+; lycopodioides, 132+; rostratum, 254+
Moenkemeyera, 253+; alomifolia, 267+; neonii, *264*; obtusifolia, *258*; uleana, *257*+; wainioi var. leuconeura, 267+, var. mollis, *257*; wainionis, *257*
Mollia cirrata, 421+
Monoclea, 244+; forsteri, 245+; gottschei subsp. elongata, 244+*, subsp. gottschei, 244+
Morales Méndez, A. *See* Marcano, V. & A. Morales Méndez, 26, 190
Mueller, D. M. J. [Review:] K. M. Van de Graaff et al. A photographic atlas for the botany laboratory, 112
Muellerella pygmaea, 156+
Mycobilimbia berengeriana, 150+; hypnorum, 150+; obscurata, 150+
Mycoblastus, 298+; alpinus, 150+
Mycocalicium subtile, 150+, 426+
Myelochroa, 426+
Myurella julacea, 132+
Myuriopsis sinica, *131*
Myurium sinicum, *131*
Myuroclada maximowiczii, 132+
Neacroporium flagelliferum, *132*
Neckera, 434+, sect. Pseudopilotrichum, *434*; subsect. Orthostichella, *434*; douglasii, 399+; extans, *173*; panduraefolia, *435*; semitorta, *436*; subdisticha, *173*; undulatifolia, 128+; yezoana, 132+
Neckeropsis, 171+; disticha, 173+, 430+; lepineana, 430+; **pocsii** sp. nov., 171*; semperiana, 276+; subdisticha, 173+
Neodolichomitra yunnanensis, 133+
Neosharpiella, 85+; aztecorum, 85+
Nephroma, 397+; arcticum, 45+, 150+; bellum, 150+, 399+; expallidum, 45+, 150+; helveticum, 399+, subsp. sipeanum, 150+; isidiosum, 150+; parile, 45+, 150+; resupinatum, 150+, 399+
Neurospora crassa, 124
Newton, A. E. [Review:] M. Mogie. The evolution of asexual reproduction in plants, 350
Nimis, P. L., J. Poelt & M. Tretiach. Caloplaca wetmorei, a new lichen species from western North America, 182
Normandina pulchella, 426+
Nostoc, 186+
Nothofagus cunninghamii, 297+; moorei, 298+
Nowellia curvifolia, 22+

O'Brien, T. J. A taxonomic note on Crassicosta hermannii, 174
Ochrolechia, 59+, 298+; androgyna, 150+; frigida, 150+; grimmiae, 150+; gyalectina, 150+; inaequatula, 150+; laevigata, 150+; oregonensis, 150+; rosella, 150+; subplicans subsp. subplicans, 150+; szatalaensis, 150+; trochophora, 409+; upsaliensis, 150+; yasudae, 409+
Octoblepharum albidum, 325
Odontoschisma macounii, 168+
Oedipodiella, 85+
Okamuraea hakoniensis, 133+
Okazaki, K. *See* Hamada, N., K. Okazaki & M. Shinozaki, 176
Omphalina hudsoniana, 143+
Omphalotus filiformis, 423+
Opegrapha stellata, 298+
Oreoweisia bogotensis, *85*+; delgadilloi, 85+; lechleri, 85+; mexicana, 85+
Orphniospora moriopsis, 150+
Orthostichella, 434+; filamentosula, 434+; **hexasticha** comb. nov., 435; **pandurifolia** comb. nov., 435; **pentasticha** comb. nov., 435; **turgidellacea** comb. nov., 435
Orthothecium hyalopiliferum, *133*; intricatum, 133+; rufescens, 276+
Orthotrichum sect. Diaphana, 405+; subg. Pulchella, 405+; acuminatum, 404+; crenulatum, 406+; diaphanum, 404+; erubescens, 133+; flowersii, 406+; fortunatii, *133*; lyellii, 399+; **macrocephalum** sp. nov., 402*; microcarpum, 408+; pallens, 407+; pumilum, 404+; tristriatum, 406+
Oxystegus cuspidatus, 133+

Pannaria ahlneri, 56+; pezizoides, 150+
Papillaria, 434; appressa, 421+; borchgrevinkii, *436*; callochlorosa, 421+; latifolia, 421+; media, 421+; nigrescens, 421+; usneoides, 421+
Parmelia, 59+; fraudans, 46+, 150+; hygrophila, 150+, 399+; omphalodes, 46+, subsp. omphalodes, 150+; salcrambidiocarpa, 298+; saxatilis, 46+, 150+; skultii, 150+; squarrosa, 150+; sulcata, 46+, 150+, 399+
Parmeliella arctophila, 150+; praetermissa, 150+; triptophylla, 150+

Parmelina galbina, 409+; metarevoluta, 409+; pseudorelicina, 298+
Parmeliopsis ambigua, 46+, 150+; hyperopta, 150+, 399+
Parmotrema conjunctum, 432+; larense, 432+; ultralucens, 432+
Pedinolejeunea uchimae, 278+
Pellia, 10+
Peltigera, 57+, 397+; aphthosa, 46+, 150+; canina, 46+, 59+, 150+; collina, 151+; degenii, 151+; didactyla, 46+, 58+, 151+; horizontalis, 151+; kristinssonii, 151+; lairdi, 440+; lepidophora, 46+, 59+, 151+; leucophlebia, 46+, 151+; malacea, 46+, 151+; membranacea, 46+, 151+; polydactyla, 46+, 151+; praetextata, 58+, 151+; retifoveata, 151+; rufescens, 46+, 151+; scabrosa, 46+, 151+; venosa, 46+, 151+
Peltula euploca, 56+
Pertusaria, 183+; alaskensis, 151+; borealis, 151+; bryontha, 151+; consocians, 426+; coriacea, 151+; dactylina, 151+; excludens, 151+; geminipara, 151+; gibberosa, 298+; gyalectina, 151+; multipuncta, 151+; oculata, 151+; panyrga, 151+; subambigens, 151+; subdactylina, 151+; subobducens, 151+; trimera, 298+; truncata, 298+; waghornei, 426+
Phacopsis huuskonenii, 156+
Phaeocalicium polyporaeum, 426+
Phaeolus schweinitzii, 322
Phaeophyscia cernohorskyi, 151+; ciliata, 151+; constipata, 46+, 151+; endococcinea, 151+; hirtella, 151+; hispidula, 151+; kairamoi, 151+; orbicularis, 58+, 151+; rubropulchra, 409+; sciastra, 46+, 58+, 151+
Phaeorrhiza nimbosa, 151+
Phascum curvicolle, 415
Philonotis bartramioides, 133+; falcata, 133+; glaucescens, 421+; mollis, 128+; nitida, 133+; revoluta, *133*; socia, *133*; thwaitesii, 133+
Phlox diffusa, 83
Phragmicoma bicolor, 423+; polycarpa, 423+
Phragmites australis, 415
Phycopeltis, 73
Phylliscum tenue, 151+
Phyllogonium viride, 421+
Physcia adscendens, 46+, 151+; aipolia, 46+, 151+; caesia, 46+, 151+; callosa, 56+; dubia, 46+, 151+; phaea, 46+, 151+; stellaris, 151+; subtilis, 151+; tenella, 151+
Physconia detersa, 151+; enteroxantha, 151+; muscigena, 46+, 151+; perisidiosa, 151+
Physolinum monilia, 186+
Picea sitchensis, 60+
Pictolejeunea picta, 325
Pilophorus cereolus, 151+
Pilosium chlorophyllum, 325
Pilotrichella, 434+; cyathipoma, 435+; filamentosula, 435+; nudiramulosa, 435+; pachygastrella, 435+; rigida, 435+; squarrulosa, 421+; turgidellacea, *435*; versicolor, 435+; viridis, 435+
Pilotrichopsis dentata, 133+
Pinnatella, 305+; alopecuroides, 307+; ambigua, 133+, 307+; amblyphylla, 307+; anacamptolepis, 307+; calcutensis, 307+; foreauana, 307+; gollanii, 307+; kuehliana, 307+; limbata, 307+; makinoa, 133+, 307+; minuta, 305+; mucronata, 307+; robusta, 307+; taiwanensis, 307+; uroclada, 307+
Pinus mugo, 158; palustris, 282
Placidiopsis cervinula, 365+; cinerascens, 365+; minor, 368+; pseudocinerea, 365+
Placodium verruculiferum, *388*
Placopsis gelida, 151+
Placynthiella icmalea 151+; oligotropha, 152+; uliginosa, 152+
Placynthium, 144+; asperellum, 152+; nigrum, 152+
Plagiobryum zierii, 133+
Plagiochasma, 250+*
Plagiochila asplenioides, 170+; corrugata, 423+; guilleminiana, 423+; heteromalla, 423+; macrifolia, 423+; rutilans, 423+; subdisticha, 423+
Plagiomnium, 10+; arbuscula, 133+; confertidens, 133+; cuspidatum, 133+, 227+; japonicum, 133+; maximoviczii, 133+; succulentum, 133+
Plagiothecium cavifolium var. cavifolium, 133+, var. fallax, 133+; denticulatum, 22+
Platismatia, 397+; glauca, 152+, 399+; herrei, 152+, 399+; lacunosa, 152+; norvegica, 152+; stenophylla, 399+
Plectocarpon lichenum, 156+
Pleuroclada albescens, 167+
Pleurozium schreberi, 127+, 358+
Pócs, T. *See* Reese, W. D., R. E. Magill & T. Pócs, 430
Poelt, J. *See* Nimis, P. L., J. Poelt & M. Tretiach, 182
Poeltinula cerebrina, 59+
Pogonatum cirratum, 133+, subsp. cirratum, 133+, subsp. fuscatum, 133+; dentatum, 133+; fastigiatum, 133+; neesii, 133+; pergranulatum, 133+; proliferum, 133+; spinulosum, 133+; subfuscatum, 133+; urnigerum, 133+
Pohlia elongata, 133+; proligera, 133+
Poli, F. *See* Gerdol, R., A. Bonora & F. Poli, 158
Polyblastia bryophila, 152+; hyperborea, 152+; obsoleta, 152+; sendtneri, 152+; terrestris, 152+; theleodes, 152+
Polychidium muscicola, 152+
Polypodium, 396
Polytrichum, 24+; aristiflorum, 421+; brasiliense, 421+; camptocaulon, 421+; commune, 133+; formosum, 133+; ohioense, 22+; strictum, 357+
Populus trichocarpa, 57
Porella navicularis, 399+
Porina, 73+; americana, 73+*; byssophila, 79+; chlorotica var. carpinea, 73+*; cinerea, 73+*; desquamescens, 76+; dolichophora, 79+; glabra, 73+; interestes, 73+*; internigrans, 73+*; luteopallens, 75+; mastodiella, 73+; mastoidea, 73+*; nucula, 75+; orientalis, 79+; subcutanea, 73+*; **subhibernica** sp. nov., 76*; subinterestes, 73+*
Porotrichum comorense, 430+
Porpidia cinereoatra, 152+; contraponenda, 152+; crustulata, 152+; flavicunda, *152*; flavocaerulescens, 152+; grisea, 152+; lowiana, 152+; macrocarpa, 152+; melinodes, 152+; speirea, 152+; superba, 152+; thomsonii, 152+; tuberculosa, 152+
Potamium commutatum, 422+
Protoblastenia calva, 152+; rupestris, 152+
Protoparmelia, 297+; badia, 152+, 297+; petraeoides, *298*; plicatula, *299*; quernea, 298+
Pseudephebe minuscula, 46+, 152+; pubescens, 46+, 152+
Pseudevernia cladonia, 426+
Pseudobarbella attenuata, 134+; laxifolia, 134+; levieri, 134+
Pseudocyphellaria, 397+; anomala, 152+, 399+; anthraspis, 399+; crocata, 152+; faveolata, 301+; granulata, 301+; rainierensis, 321+
Pseudoleskeopsis japonica, *134*; zippelii, 134+
Pseudopilotrichum, *434*+
Pseudosymblepharis angustata, 134+
Pseudotaxiphyllum pohliaecarpum, 134+, 276+
Pseudotsuga menziesii, 321+, 396+

Psilolechia lucida, 152+
Psora cerebriformis, 152+; decipiens, 152+; himalayana, 152+; lurida, 152+; montana, 56+; nipponica, 59+; plicatula, *299*; rubiformis, 59+, 152+
Psoroma hypnorum, 46+, 152+
Psorotichia schaereri, 152+
Pterobryella longifrons, 227+; papuense, 227+; vagapensis, 227+
Pterygoneurum, 412+; arcticum, *412*; californicum, 412+; **compactum** sp. nov., 412*; crossidioides, 412+; kemsleyi, 412+; koslovii, 412+; lamellatum, 412+; macleanum, 412+; medium, 412+; ovatum, 412+; sampaianum, 412+; smardeanum, *412*; subsessile, 412+
Ptilidium ciliare, 168+; pulcherrimum, 22+, 170+
Ptychomitrium, 34+; angustifolium, *34*; gardneri, 37+, 134+; kiusiuense, *37*; leibergii, *34*; longisetum, *37*; microcarpum, *34*, var. microcarpum, *34*; polyphylloides, *37*, *134*; polyphyllum, 37+; robustum, *37*; sinense var. microcarpum, *34*; standleyi, 37+; viride, *37*; wilsonii, 134+
Ptychomnion aciculare, 227+; cygnisetum, 227+
PURSELL, R. A. Taxonomic notes on neotropical Fissidens, 253
Pyrenopsis polycocca, 152+
Pyrenula laevigata, 426+
Pyrrhobryum bifarium, 227+; dozyanum, 134+, 227+; spiniforme, 227+, 263+
Pyrrhospora, 296+; elabens, 296+; laeta, 297+; petraeoides, *298*; sanguinolenta, 297+
Pyxine, 432+

Quercus douglasii, 366; faginea, 405+; garryana, 40; ilex, 405+; laevis, 282; lobata, 366; pyrenaica, 405+

Racodium, 186+; rupestre, 186+
Racomitrium anomodontoides, 134+; fasciculare var. atroviride, *134*; himalayanum, 134+; subsecundum, 134+
Racopilum, 10+; aristatum, 134+, 227+; cuspidigerum, 10+; ferriei, 134+; spectabile, 10+, 227+; strumiferum, 227+
Radula acuminata, 279+; kojana, 279+; macrostachya, 325; oyamensis, 279+; tomentosum, 422+; uvifera, 314+; xalapensis, 423+
Ramalina, 26+; africana, 29+, 190+*; **andina** sp. nov., 26*; **chiguarensis** sp. nov., 27*; **crispata** sp. nov., 28*; cuspidata, 29+; decipiens, 29+; farinacea, 152+, 399+; hamulosa, 27+; intermedia, 426+; longissima, 27+; **lopezii** sp. nov., 29*; **microphylla** sp. nov., 30*; nodosa, 31+; obtusata, 152+; pacifica, 26+; pollinaria, 152+, 426+; pusiola, 26+; **santanensis** sp. nov., 31*; siliquosa, 29+; subbreviuscula, 176+; subcomplanata, 178+; sublitoralis, 176+; **tenuissima** sp. nov., 31*; thrausta, 152+, 426+; **tovarensis** sp. nov., 33*, 190+; usnea, 26+
Ramboldia gen. nov., 296; **brunneocarpa** sp. nov., 297*; **petraeoides** comb. nov., 298*; **plicatula** comb. nov., 299*; **stuartii** comb. nov., 299*; **subnexa** comb. nov., 301*
Ramularia peltigericola, *57*
REDFEARN, P. L., JR., [Review:] J. Beever et al. The mosses of New Zealand, 110
REDFEARN, P. L., JR., B. ALLEN & S. HE. New distributional records for Chinese mosses, 275
REESE, W. D. A new name and a new species in Syrrhopodon subgenus Pseudo-calymperes from Cameroon, 416
REESE, W. D. Per Dusén's African species of Syrrhopodon subgenus Pseudo-calymperes, 371
REESE, W. D. [Review:] R. C. Lobato Lisboa. Musgos Acrocárpicos do Estado de Rondônia, 467
REESE, W. D. [Review:] S. Metzler & V. Metzler. Texas mushrooms. A field guide, 222
REESE, W. D. [Review:] R. H. Zander. Genera of the Pottiaceae: mosses of harsh environments, 216
REESE, W. D., R. E. MAGILL & T. PÓCS. Mitthyridium micro-undulatum subsp. comoroensis subsp. nov. from Mayotte, 430
REESE, W. D. & H. STREIMANN. Calymperes subserratum (Musci), new to eastern Malesia, with notes on C. serratum and C. subulatum, 80
Rhabdoweisia crispata, 134+; denticulata, *134*; sinensis, *134*
Rhacocarpus humboldtii, 422+; inermis, 422+; piliformis, 422+
Rhaphidostegium cyparissoides, 422+; kegelii, 422+; pseudo-callidioides, 422+
Rhizocarpon alpicola, 152+; badioatrum, 152+; chioneum, 152+; cinereonigrum, 152+; copelandii, 153+; disporum, 153+; eupetraeoides, 153+; eupetraeum, 153+; geminatum, 153+; geographicum, 153+; grande, 153+; hochstetteri, 153+; inarense, 153+; polycarpum, 153+; rittokense, 153+; superficiale, 153+; cf. umbilicatum fo. pseudospireum, 153+
Rhizogonium distichum, 227+; novae-hollandiae, 227+; spiniforme, 422+
Rhizomnium horikawaea, 134+
Rhizoplaca chrysoleuca, 153+; melanopthalma, 153+
Rhodobryum giganteum, 134+
Rhytidiadelphus japonicus, 134+; subpinnatus, 134+
Riccardia graeffei, 87+
Riccia dictyospora, 10
Rigodium, 225+
Rimelia **bonplandii** sp. nov., 432*; diffractaica, 432+; subisidiosa, 433+
Rinodina adirondackii, 426+; archaea, 153+; ascociscana, 426+; bischoffii, 153+; darrovii, 439+; laevigata, 153+; occidentalis, 153+; oregona, 153+; pyrina, 153+; roscida, 153+; turfacea, 153+
ROOKS, P. E. *See* SHAW, A. J. & P. E. ROOKS, 1
Ropalospora lugubris, 153+
ROS, R. M. *See* CANO, M. J., J. GUERRA & R. M. ROS, 412
ROSENTRETER, R. *See* GOWARD, T., P. DIEDERICH & R. ROSENTRETER, 56
RUSHING, A. E. [Review:] M. R. Crosby & R. E. Magill. (Compilers.) Index of mosses, 1990–1992, 474
RUSHING, A. E. [Review:] N. G. Miller. (ed.) Bryophyte systematics, 465

Saccharomyces cerevisiae, 118+
Saccobasis, 166+; polita, 166+
Sagiolechia rhexoblephara, 153+
Sakuraia conchophylla, 134+
SALAZAR ALLEN, N. Syrrhopodon simmondsii new to Panama and Central America, 325
Salix, 59; nivalis, 83
Sarcogyne clavus, 153+; regularis, 153+
Sarracenia, 280
Sarrameana tasmanica, 298+
Scapania, 252+; americana, 168+; bolanderi, 170+, 399+; mucronata, 170+; umbrosa, 170+
Schaereria tenebrosa, 153+
Schistidium apocarpum, 134+; confertum, 134+; gracile, *134*; liliputanum, 128+; strictum, 134+
Schizymenium, 1
Schlotheimia appressifolia, 227+; campylopus, 422+; fusco-viridis, 422+; pungens, 134+; robusticuspis,

227+; serricalyx, 227+; tecta, 422+; torquata, 227+; trichomitria, 422+; wainioi, 422+
Schoenobryum coffeae, 89+; concavifolium, 89+; gardneri, 89+
SCHUSTER, R. M. [Review:] W. N. Stewart & G. W. Rothwell. Paleobotany and the evolution of plants, 463
SCHUSTER, R. M. *See* ENGEL, J. J. & R. M. SCHUSTER, 63
Schwetschkea gymnostoma, 134+
Scirpus cespitosus, 284
Sclerotinia trifoliorum, 124
Scopelophila cataractae, 47+
Scutula miliaris, 56+
SELVA, S. B. Lichen diversity and stand continuity in the northern hardwoods and spruce–fir forests of northern New England and western New Brunswick, 424
Sematophyllum argyro-viride, 422+; subhumile, 134+; subpungifolium, 422+
Sendtnera juniperina, 423+
SEPPELT, R. D. *See* MELICK, D. R. & R. D. SEPPELT, 13
SHAW, A. J. [Review:] H. C. Greven. Changes in the Dutch bryophyte flora and air pollution, 220
SHAW, A. J. Systematics of Mielichhoferia (Bryaceae: Musci) II. Morphological variation among disjunct populations of M. elongata and M. mielichhoferiana, 47
SHAW, A. J. & P. E. ROOKS. Systematics of Mielichhoferia (Bryaceae: Musci) I. Morphological and genetic analyses of M. elongata and M. mielichhoferiana, 1
SHINOZAKI, M. *See* HAMADA, N., K. OKAZAKI & M. SHINOZAKI, 176
Silene acaulis, 83
SILLETT, S. C. Growth rates of two epiphytic cyanolichen species at the edge and in the interior of a 700-year-old Douglas fir forest in the western Cascades of Oregon, 321
Simplicidens, 253+; andicola, *255*; plumula, *255*
Solidago multiradiata, 83
Solmsiella biseriata, *131*
Solorina bispora, 46+, 153+; crocea, 46+, 153+; saccata, 46+, 153+; spongiosa, 153+
Solorinella asteriscus, 59+
Sphaerellothecium araneosum, 56+
Sphaerophorus fragilis, 46+, 153+; globosus, 46+, 153+, 399+; melanocarpus, 153+; tuckermanii, 153+
Sphagnum, 10+, 158+, 252+, 284+; angustifolium, 284+, 357+; balticum, 284+; bartlettianum, 280+; capillifolium, 158+, 280+; compactum, 280+, 284+; cuspidatum, 280+, 284+; cyclophyllum, 280+; fallax, 158+, 284+; fimbriatum, 282+; flexuosum, 285+; fuscum 17+, 285+; girgensohnii, 282+, 286+; henryense, 280+; junghuhnianum, 134+; lindbergii, 284+; macrophyllum, 280+; magellanicum, 158+, 284+, 357+; majus, 284+; molle, 280+; nemoreum, 285+; ovatum, 134+; palustre, 285+; papillosum, 160+, 284+; perichaetiale, 280+; quinquefarium, 282+; riparium, 284+; rubellum, 284+; russowii, 282+, 284+; squarrosum, 284+; strictum, 280+; subsecundum var. inundatum, 284+; tenellum, 284+; tenerum, 280+, var. tenerum, 280+, var. **virginicum** var. nov., 282*
Spinacia oleracea, 159
Spiridens camusii, 227+; vieillardii, 227+
Sporastatia testudinea, 153+
Squamarina lentigera, 153+
Squamidium, 434+
STANISLAUS, R. C. & N. C. MARAVOLO. The influence of polyamines on senescence in Marchantia polymorpha, 162
Staurothele areolata, 153+, 368+; clopima, 368+; drummondii, 153+; elenkinii, 153+; fissa, 153+
Steereomitrium minutum, 63+
Stenocybe major, 426+
Stereocaulon alpinum, 46+, 153+; arenarium, 153+; botryosum, 153+; capitellatum, 153+; condensatum, 153+; depreaultii, 409+; glareosum, 153+; grande, 153+; incrustatum, 153+; intermedium, 153+; paschale, 46+, 153+; rivulorum, 46+, 153+; saxatile, 154+; subcoralloides, 154+; tennesseense, 409+; tomentosum, 46+, var. alpestre, 154+, var. tomentosum 154+; vesuvianum, 154+
Sticta arctica, 154+; fuliginosa, 154+; wrightii, 56+
STOTLER, R. [Review:] H. Bischler-Causse. Marchantia L. The European and African taxa, 351
Strangospora microhaema, 426+
STREIMANN, H. *See* REESE, W. D. & H. STREIMANN, 80
STUCKEY, R. L. Emanuel David Rudolph (1927–1992): Polar lichenologist and historian of botany, 437
STUCKEY, R. A. *See* BURK, W. R. & R. A. STUCKEY, 443
Suaeda vera, 415
Symblepharis breviseta, *128*; **guizhouensis**, 130+, nom. nov., 134; reinwardtii, 134+
Symphyodon echinatus, 276+; pygmaeus, 276+
Synechococcus, 186+
Syrrhopodon subg. Pseudo-calymperes, 371+, 416+; subg. Calymperopsis, *416*; africanus, 371+*, 416+, 431+; albovaginatus, 80+; argenteus, 422+; capillaceus, 422+; carassensis, 422+; "cuspidato," *431*; cymbifolius, 419+; disciformis, 373+; duseni, *373*; gardneri, 134+; gaudichaudii, 325; gracilescens, 422+; japonicus, 134+; martinicensis, *374*; muelleri, 80+; parasiticus, 263+, 325, 371+, 419+, var. disciformis, 431+; **perdusenii**, 371+, nom. nov., 416*; simmondsii, 325; spurio-disciformis, *371*+; subdisciformis, 371+, *416*; "subflavo," *431*; **virgulicola**, 371+, sp. nov., 419*; vitiense, *81*; wainioi, 422+

Taiwanobryum speciosum, 135+
Takakia, 244+
TAN, B. C., QI-WEI LIN, M. R. CROSBY & PAN-CHENG WU. A report on the 1991 Sino–American bryological expedition to Guizhou Province, China: New and noteworthy additions of Chinese moss taxa, 127
Taxiphyllum subarcuatum, 128+, 276+
Taxithelium planum, 325
Taxodium, 89
Taxus brevifolia, 322
Telaranea nematodes, 71+*
Tephromela aglaea, 154+; armeniaca, 154+; atra, 154+, 301+
Tetraphis pellucida, 20+; 135+
Thamniopsis utacamundiana, *131*
Thamnium laevinervis, 135+
Thamnobryum cf. alopecurum, 135+, 227+; plicatulum, 135+; sandei, *135*; subserratum, 135+
Thamnolecania macquariensis, 440+
Thamnolia subuliformis, 46+, 83+, 154+; vermicularis, 17+, 46+, 154+
Thelidium absconditum, 154+
THIERS, B. M. [Review:] D. Sharma & S. C. Srivastava. Indian Lepidozineae (a taxonomic revision), 348
THIERS, B. M. & W. R. BUCK. Recent literature on hepatics and anthocerotes—135, 91;—137, 326
THOMSON, J. W. [Review:] J. Poelt & E. Hinteregger. Beiträge zur Kenntnis der Flechtenflora des Himalaya. VII. Die Gattungen Caloplaca, Fulgensia, und Ioplaca (mit englischem Bestimmungsschlüssel), 350
THOMSON, J. W. & T. AHTI. Lichens collected on an Alaska Highway expedition in Alaska and Canada, 138
Thrombium epigaeum, 154+
Thuidium cymbifolium, 135+; kanedae, 135+; philiber-

tii, 135+; pinnatulum, 422+; subdelicatulum, 422+; subglaucinum, 135+
Thuja plicata, 322
Thysanomitrium carassense, 422+; richardii, 422+
Timmia austriaca, 227+; bavarica, 227+; megapolitana, 227+
Timmiella anomala, 135+
Toloxis nom. nov., 436; **borchgrevinkii** comb. nov., 435; **imponderosa** comb. nov., 436; **semitorta** comb. nov., 436
Toninia caeruleonigricans, *154*; candida, 59+; sedifolia, 154+; tristis, 60+, 154+, subsp. asiae-centralis, 60+
Tortula brevissima, 415; laevipila, 404+; vahliana, 415
"Torula lichenum," 156+
Trachelospermum jasminoides, 277+
Trachycystis microphylla, 135+, 227+; ussuriensis, 129+, 135+
Trachypodopsis humilis, 135+; serrulata var. crispatula, 135+
Trapelia corticola, 56+
Trapeliopsis granulosa, 154+; viridescens, 60+
Trebouxia erici, 118+
Trematodon reflexus, 422+
Tremolecia atrata, 154+; jurana, 154+
Trentepohlia, 73, 186+
TRETIACH, M. *See* NIMIS, P. L., J. POELT & M. TRETIACH, 182
Treubia lacunosa, 65+; tasmanica, 64+
Trichostomum brachydontium, 135+
Trichothelium alboatrum, 272+; album, 272+; **marianense** sp. nov., 272*; minus, 274+
Trilophozia, 166+
Tritomaria sect. Tritomaria, 166+; subg. Saccobasis, 166+, subg. Tritomaria, 166+; exsecta, 166+; exsectiformis, 166+; polita, 166+; quinquedentata var. quinquedentata, 166+, var. turgida, 166+; scitula, 166+; trilophozia, 166+
Trocholejeunea sandvicensis, 279+
Tsuga heterophylla, 59, 322, 396+; plicata, 396+
Tuckermannopsis americana, 154+; aurescens, 426+; chlorophylla, 154+; merrillii, 154+; orbata, 154+; platyphylla, 154+; sepincola, 154+

Ulota crispa, 399+; megalospora, 399+
Umbilicaria angulata, 154+; aprina, 60+, var. halei, *60*; arctica, 46+, var. diomedensis, 154+; cinereorufescens, 60+; cylindrica, 154+; decussata, 14+; deusta, 46+, 154+; havaasii, 154+; hyperborea, 46+, 154+; lyngei, 46+; muehlenbergii, 46+; polyphylla, 46+, 154+; proboscidea, 46+, 154+; torrefacta, 46+, 154+; vellea, 46+, 60+, 154+
UPGMA, 48+
UPRETI, D. K. Notes on corticolous and saxicolous species of Porina from India, with Porina subhibernica sp. nov., 73
Usnea, 397+; angulata, 154+; capitata, 154+; cavernosa, 154+; ceratina, 154+; filipendula, 154+; inermis, 301+; lapponica 46+, 57+, 426+; longissima, 154+, 426+; mutabilis, 409+; occidentalis, 154+; oncodes, 298+; plicata, 399+; rubicunda, 426+; scabrata, 399+, var. nylanderiana, 154+; sphacelata, 14+; subfloridana, 154+; substerilis, 154+; wasmuthii, 154+; wirthii, 56+

Varicellaria rhodocarpa, 154+
Verrucaria, 377+; arctica, 154+; compacta, 369+; devergens, 154+; deversa, 154+; funckii, 154+; **lobata** sp. nov., 155*; margacea, 155+; maura, 155+; nigrescens, 155+; rupestris, 155+; viridula, 155+; zamenhofiana, 365+
Vestergrenopsis elaeina, 56+, 155+; isidiata, 155+
VISNADI, S. R. *See* VITAL, D. M. & S. R. VISNADI, 71
VITAL, D. M. & S. R. VISNADI. New records and notes on Brazilian hepaticopsida, 71
VITT, D. H. *See* CAO, TONG & D. H. VITT, 34
VITT, D. H. *See* LI, YENHUNG & D. H. VITT, 357
VITT, D. H. [Review:] J. M. Glime. The elfin world of mosses and liverworts of Michigan's upper peninsula and Isle Royale, 472
VITT, D. H. & P. KUHRY [Review:] J. T. A. Verhoeven. (ed.) Fens and bogs in the Netherlands: Vegetation, history, nutrient dynamics and conservation, 218
Vulpicida canadensis, 155+; pinastri, 42+, 155+; tilesii, 46+, 83+, 156+

Weissia controversa, 135+; edentula, 135+; longidens, 135+
Weymouthia, 434+
WOJTUŃ, B. Element contents of Sphagnum mosses of peat bogs of lower Silesia (Poland), 284
WU, PAN-CHENG *See* TAN, B. C., QI-WEI LIN, M. R. CROSBY & PAN-CHENG WU, 127

Xanthoparmelia coloradoensis, 59+; cumberlandia, 59+; planilobata, 56+; somloensis, 156+; taractica, 46+; wyomingica, 59+
Xanthopyrenia heardensis, 440+
Xanthoria candelaria, 46+, 156+; elegans, 46+, 156+, 388; fallax, 46+, 156+; polycarpa, 156+; sorediata, 156+
Xenochila, 65+
Xenothallus, 64+
Xylaria magnoliae, 124
Xylographa abietina, 156+

Yarrowia lipolytica, 124
YE, LI-XIN *See* ZHU, RUI-LIANG, LI-XIN YE & HUAN-ZHONG CAI, 277

ZHU, RUI-LIANG, LI-XIN YE & HUAN-ZHONG CAI. Epiphyllous liverworts of Fengyangshan nature reserve, Zhejiang Province, China, 277
Zoopsis antillana, 71+